

# Toxic Interactions among Environmental Pollutants: Corroborating Laboratory Observations with Human Experience

Kannan Krishnan and Jules Brodeur

Département de médecine du travail et d'hygiène du milieu, Faculté de médecine, Université de Montréal, Montréal, Canada

Combined exposures to multiple chemicals may result in interactions leading to a significant increase or decrease in the overall toxicity of the mixture compared to the summation of the toxicity of the components. A large number of chemical interactions have been described in animal studies by administering high doses of chemicals by routes and scenarios often different from anticipated human exposures. Though limited, there is some evidence for the occurrence of several supra-additive (the combined effects are greater than the simple summation of the individual effects) and infra-additive (the combined effects are smaller than the simple summation of the individual effects) chemical interactions in humans. For example, toxicokinetic interactions between several solvents have been found to occur in the workplace, whereas those involving pesticides have been reported less frequently, especially during accidental exposures. Toxic interactions involving nutritionally important metals and metalloids appear to occur more frequently, since several of them have an important role in a variety of physiological and biochemical processes. On the contrary, there is not much evidence to confirm the occurrence of toxic interactions among the commonly encountered inorganic gaseous pollutants in humans. Overall, the majority of chemical interactions observed in animal studies have neither been investigated in humans nor been extrapolated to humans based on appropriate mechanistic considerations. Future research efforts in the chemical interactions arena should address these issues by focusing on the development of mechanistically and biologically based models that allow predictions of the extent of interactions likely to be observed in humans.—*Environ Health Perspect* 102(Suppl 9):11–17 (1994)

Key words: toxic interactions, chemical interactions, potentiation, synergism, antagonism

## Introduction

Toxic interaction refers to the qualitative and/or quantitative modification of the toxicity of one chemical by another, the process principally occurring within the organism after the exposure phase (1). Interactions can either result in greater-than-additive or less-than-additive toxic response. Over a thousand studies published to date report the occurrence of supra- or infra-additive toxicity from combined exposure to two chemicals (2). The interactive toxicity resulting from combined exposures to chemicals is a consequence of the alteration of the toxicokinetics and/or toxicodynamics of one chemical by another. Interference at the kinetic level would imply a modulation of absorption, distribution, metabolism and/or excretion of one chemical by another. Interference at the toxicodynamic level might involve a competition between two chemicals for binding to

the target site or an alteration by one chemical of the susceptibility of the target cells to the effects of another agent.

Despite the report of the occurrence of several significant interactions between environmental pollutants, their relevance for humans and relative importance for regulation remain unclear and ill-defined. This situation is a consequence of the lack of chronic exposure studies with chemical mixtures and the lack of understanding of the mechanistic basis of interactions at a quantitative level. Until these issues are resolved, we probably would not be able to confidently use animal data on interactions to make quantitative predictions for humans. However, there is some direct and/or epidemiological evidence for the occurrence of several supra-additive and infra-additive chemical interactions in humans. In this article, we corroborate laboratory observations with human experience as they relate to toxic interactions among environmental pollutants and propose an approach to consider data on toxic interactions for human health risk assessment.

### Toxic Interactions among Gaseous Pollutants

Interactions among gaseous air pollutants most commonly involve physicochemical mechanisms rather than toxicodynamic/kinetic inter-


ferences. A number of animal studies have suggested the occurrence of supra-additive and infra-additive interactions among gaseous pollutants (Figures 1,2). Of those interactions observed in laboratory studies, there is evidence suggesting that a few of them may have actually been experienced by humans. For example, the antagonistic interaction resulting from combined exposure to sulfur dioxide and ammoniacal compounds is thought to have been encountered during the London fog disaster of 1952 (3). The supra-additive toxicity resulting from combined exposures to sulfur dioxide and ozone may have been a causative factor of the high mortality of Japanese children observed in the early seventies (4). The reported higher incidence of cancer among workers in arsenic-smelting facilities has been explained by the supra-additive effects between arsenic and benzo[a]pyrene observed in laboratory animals (5). However, the majority of the interactions involving commonly occurring gaseous pollutants (i.e., NO<sub>x</sub>, SO<sub>x</sub>, ozone, aerosols) that have been well characterized in animal studies do not seem to occur in humans (6–11).

### Toxic Interactions among Pesticidal Chemicals


Humans ingesting food preparations contaminated with one or more pesticides, workers in pesticide manufacturing and

This article was presented at the IV European ISSX Meeting on Toxicological Evaluation of Chemical Interactions: Relevance of Social, Environmental and Occupational Factors held 3–6 July 1992 in Bologna, Italy.

Address correspondence to Dr. K. Krishnan, Département de Médecine du Travail et d'Hygiène du Milieu, Faculté de Médecine, Université de Montréal, Case Postale 6128, Succursale A, Montréal, P.Q., Canada, H3C 3J7. Telephone (514) 343-6581. Fax (514) 343-2200.



**Figure 1.** Supra-additive interactions among gases and particulates demonstrated in animal studies (2).



**Figure 2.** Infra-additive interactions among gases and particulates demonstrated in animal studies (2).

packaging units, and agricultural workers who prepare, mix, and apply pesticides in the fields are all potentially exposed to more than one pesticide on the same or successive days. Interactions among pesticides often involve either an induction or an inhibition of a particular metabolic pathway.

The supra-additive and infra-additive binary interactions reported among commercially important pesticidal chemicals in animal studies are summarized in Figures 3 and 4. The only instance where the occurrence of an interaction has been suspected pertains to a poisoning episode in 1976, when hundreds of Pakistani spraymen became ill following the use of a water-dispersible formulation of malathion (12). In this instance, isomalathion, an impurity of malathion preparation, formed in great quantities under improper storage conditions, accounted for the enhanced cholinergic effects. Animal studies have shown that isomalathion is more potent than malathion as an inhibitor of acetylcholinesterase and also potentiates malathion toxicity by interfering with the carboxylesterase pathway of malathion detoxication (13,14).

### Toxic Interactions among Metals/Metalloids

Humans are exposed to low concentrations of several metals and metalloids arising from fossil fuel burning, vehicular exhausts, metallic ore smelting, and waste incineration. Occupational exposures may result in substantial body burden of metals

depending upon the type of operation and facility. With metals and metalloids, toxicological interactions occur at both extremes of tissue exposure, i.e., deficiency and excess. This property is a consequence of several metals being required for a variety of physiological processes, e.g., as cofactors of enzymes.


The supra-additive and infra-additive interactions among metals and metalloids described in animal studies are depicted in Figures 5 and 6. Of these, the following infra-additive interactions have been investigated in humans: selenium and mercury, zinc and lead, and iron and mercury, cadmium or lead.

Selenium-induced protection of mercury toxicity in rats has been reported widely in the literature (15–17). There is at least one case report of beneficial effect due to selenium-mercury interaction in humans (18). These authors reported that allergic reactions, skin rash, incoordination and memory loss in a 39-year-old woman sensitive to mercury-based paints and sprays, were markedly reduced after she took periodic selenium supplements.


Suzuki et al. (19), measured the concentrations of trace elements in foodstuffs from a Japanese area with elevated intake of methyl mercury and found comparable levels of selenium as well. These authors considered that the alleviating effect of selenium is the probable reason for the toxic effects of mercury not being evident in that population. Some autopsy studies have reported that both mercury and selenium accumulated in tissues at an approximate molar ratio of 1:1 suggesting a natural protective effect by selenium of mercury toxicity (20,21).

Regarding the antagonistic interaction between mercury and iron, there is one corroborative accidental report (22). A 5-year-old boy who ingested an open battery containing iron and mercuric oxide experienced no significant uptake of mercury. This has been attributed to the reduction of  $\text{HgO}$  into insoluble elemental mercury by iron present at the gastric site.

Animal studies have indicated that supplemental uptake of iron decreases the absorption of lead, antagonizing lead-induced anemia; conversely, prevalence of iron deficiency has been associated with increasing severity of lead poisoning (23–28). It has therefore been suggested that the correction of iron deficiency among human populations can generally reduce lead toxicity (29). Interestingly, a negative correlation between blood lead levels and body burden of iron in human



**Figure 3.** Supra-additive interactions among pesticides demonstrated in animal studies (2). Abbreviations: BPMC, 2-sec-butylphenylmethylcarbamate; DDT, 1,1,1-trichloro-2,2-bis-(4-chlorophenyl) ethane.




**Figure 4.** Infra-additive interactions among pesticides demonstrated in animal studies (2).


populations has been reported by several authors (30–34).

Similar protective effects of zinc on lead toxicity have been reported in certain animal and human studies. In a survey of industrial workers, Dukiewicz et al. (35) found significantly reduced excretion of aminolevulinic acid in workers exposed to both lead and zinc compared to those exposed to lead alone. Though the protective action of zinc in the case of lead poisoning is suggested (36), its relevance for humans still remains uncertain (37,38).


The absorption of zinc is reduced in the presence of inorganic iron as a result of a competition for intestinal carrier sites (39). According to this mechanistic description, 25 mg of total ions (as the sum of zinc and




**Figure 5.** Supra-additive interactions among metals and metalloids demonstrated in animal studies (2).



**Figure 6.** Infra-additive interactions among metals and metalloids demonstrated in animal studies (2).



**Figure 7.** Infra-additive and inhibitory interactions among solvents demonstrated in animal studies (2).



**Figure 8.** Supra-additive interactions among solvents demonstrated in animal studies (2).

iron administered orally) is the point of saturation, at which competitive effect will begin to be expressed (39). Evidence in support of interaction between iron and zinc/cadmium has been obtained in human studies (40).

### Toxic Interactions among Solvents

Combined exposure to organic solvents often results in a mutual inhibition of their metabolism, since most of them appear to be metabolized by cytochrome P450 2E1 at low exposure concentrations. Figure 7 presents the inhibitory interactions among solvents determined in laboratory animal studies. Of the metabolic inhibition interactions observed in animal studies, the following have been confirmed in human volunteer/occupational exposure studies: trichloroethylene and 1,1,1-trichloroethane (41), benzene and toluene (42), ethylbenzene and *m*-xylene (43), xylene and toluene (44), trichloroethylene and tetrachloroethylene (45), *m*-xylene and methyl ethyl ketone (46) and *m*-xylene and isobutanol (47). These inhibitory metabolic interactions are characterized by reduced/delayed production and excretion of metabolites, and/or increased concentrations of parent chemical(s) in the blood and expired air. However, the health significance of several of these metabolic interactions has not been confirmed in workers occupationally exposed to binary solvent mixtures (48,49). The inhibitory metabolic interactions can be expected to result in supra-additivity with respect to the toxicity of the parent chemicals.

Supra-additive toxicity during exposure to binary mixtures of solvents has often been observed in animal studies, whenever one of the two solvents was a potent inducer of activating enzymes, or an inhibitor of detoxication enzyme system (50,51). Thus, most of the solvents that induce P4502E1 upon prior administration (e.g., ketones) have been shown to potentiate the toxicity of other solvents bioactivated by the same isoenzyme (Figure 8).

In human experience, isopropanol, a ketogenic chemical, was the suspect potentiator of the hepato-renal toxicity of carbon tetrachloride in two separate industrial accidents (52,53). Similarly, methyl ethyl ketone-induced potentiation of *n*-hexane/MnBK neurotoxicity, through the enhanced formation of 2,5-hexanedione, appears to have been responsible for the outbreak of an occupational neuropathy among the textile workers in Ohio (54,55) and among glue sniffers in West Berlin in the seventies (56).

### Relevance of Animal Data on Chemical Interactions for Quantitative Risk Assessment

In evaluating the relevance of chemical interactions for regulatory purposes, it is important to consider the underlying mechanisms. Understanding the mechanistic basis enables us to determine whether or not an interaction will occur at low exposure levels in animals, and if it will occur at all in other species, especially, humans. Table 1 presents some of the mechanisms of interaction unequivocally demonstrated in animal and/or human studies.

The data generated in most interaction studies are qualitative, supporting or suggesting a specific mechanism as the possible cause of the infra- and supra-additivity. For quantitative risk assessment (QRA) purposes, we need quantitative mechanistic data on chemical interactions. For example, what is the dose-response for the induction or inhibition of metabolism of one chemical by another? What is the quantitative relationship between two chemicals competing for protein binding? Such quantitative analyses of chemical interactions can be conducted when the interrelationships among the critical determinants of chemical disposition, action and interaction are identified and integrated within a biologically based modeling framework. To date, this quantitative modeling approach has been applied only for a few binary chemical mixtures (72).

Since the quantitative mechanistic basis of chemical interactions has not been elucidated for majority of interacting chemical

combinations, their relevance for humans cannot be evaluated confidently. On the other hand, regardless of the level of mechanistic understanding of interchemical interactions, there are instances that require the evaluation of the relevance of data on chemical interactions for specific purposes (e.g., for ensuring worker safety).

Under these circumstances, we are obliged to analyze the available data and come up with a tentative conclusion regarding the importance of an interaction for a given exposure situation. There are certain types of data which suggest that a particular interaction is unimportant for humans. These relate primarily to instances where the animal studies show the existence of thresholds for interactions which exceed the maximum allowable exposure concentrations for humans, and where animal studies confirm the occurrence of interactions which do not occur in humans exposed to low levels of both chemicals.

An example of the former case is the EPN (*O*-ethyl-*O*-4-nitrophenyl phenyl phosphonothioate) malathion potentiation. This interaction, shown to be important at high doses (73), was not apparent when animals were fed recommended daily tolerance levels of both chemicals (74), suggesting that the potentiation may not occur in humans exposed to much lower concentrations of these chemicals. Thus, Moeller and Rider (75), giving small oral doses of EPN and malathion in combination to human volunteers, found no evidence of EPN potentiation of malathion toxicity.

An example of the second type is the interaction among several commonly occurring air pollutants. Pulmonary biochemical and morphological changes induced by nitrogen dioxide have been reported to be enhanced by coexposure to acidic aerosols (76). This interaction occurs even at levels of 1 mg/m<sup>3</sup> of sulfuric acid aerosol and 2 ppm of nitrogen dioxide. But in human exposure studies, Stacey et al. (7) could not detect any synergistic effect in people exposed to low levels of both chemicals. Similarly, synergistic interaction between ozone and sulfuric acid aerosols has been demonstrated in animals exposed to atmospheric concentrations of 0.12 ppm and 5 mg/m<sup>3</sup>, respectively (77). Yet, Horvath et al. (10) found that a 2-hr exposure to a mixture of 0.2 ppm of ozone and 1.2 to 1.6 mg/m<sup>3</sup> of sulfuric acid aerosol did not induce effects beyond what could be attributed to human exposure to ozone alone.

Detailed investigations of this kind have not been conducted for the majority

of the several hundred interactions reported to occur among environmental pollutants at high doses (2). Neither chronic animal exposure data nor human volunteer exposure data are available for these binary mixtures, rendering the assessment of their relevance for humans difficult. When confronted with a particular exposure situation involving chemical interactions, one should evaluate the available information on that interaction (or for a similar chemical combination), on the basis of factors such as dose administered


and species used, to determine the relevance for humans (Figure 9). Accordingly, if such an evaluation indicates a high probability for occurrence of an interaction between two chemicals encountered in a particular workplace setting, then necessary precautions should be taken to avoid such a combined exposure. In the following paragraphs, we discuss four categories of data that should be given consideration in decreasing order of importance, while evaluating the relevance of information on chemical interactions for humans.

### Interactions Demonstrated in Humans Exposed At or Below Allowable Exposure Concentrations (TLV, RfC, RfD)

Toxic interactions demonstrated in humans, in most cases, pertain to occupational exposures involving solvents more frequently than other types of contaminants. Toxicokinetic interactions among various solvents often arise from the mutual inhibition of their metabolism, thus resulting in increased blood levels. Such inhibitory metabolic interactions may

**Table 1.** Examples of mechanisms of toxicokinetic interactions among chemicals.

Basis of interaction	Interacting chemicals	Mechanism of interactive effects	References	
Absorption				
	Percutaneous	<i>m</i> -Xylene and isobutanol	Reduced absorption of both compounds due to dehydration of skin elicited by isobutanol	Rilhimaki (47)
		Dimethyl sulfoxide (DMSO) and pesticides	Increased dermal absorption of pesticides and other chemicals when they are mixed with DMSO, which disrupts the cellular permeability and acts as a "penetrant carrier"	Jacob et al. (57) Hayes and Pearce (58)
Pulmonary	Hydrogen cyanide (HCN) and carbon monoxide	Increased pulmonary uptake of air contaminants when they are present along with HCN, which at low concentrations increases the pulmonary ventilation rate	Dreisbach (59)	
Gastrointestinal	Lead and iron	Decreased absorption of lead in the presence of iron due to competition for transport sites in the intestinal mucosa.	Conrad and Barton (60)	
Distribution				
Tissue distribution	Lead and dithiocarbamates	The lipophilic lead-dithiocarbamate complex has a greater capacity than lead alone to penetrate the blood-brain barrier and causes a greater accumulation in the lipid-rich brain components.	Osakrsson (61)	
Protein binding	Organochlorine and organophosphate pesticides	Organochlorine pesticides not only enhance the biotransformation of organophosphates, but also enhance their binding to plasma proteins and nonspecific esterases.	Trioloand Coon (62) Cohen and Murphy (63)	
Metabolism				
Phase I				
Induction	Methyl <i>n</i> -butyl ketone (MnBK) and chloroform	Increased bioactivation of chloroform due to induction of hepatic microsomal P450 2E1 by MnBK.	Brady et al. (51)	
Inhibition	Dithiocarbamates and chloroform	Decreased bioactivation of chloroform due to inhibition of hepatic microsomal enzymes by dithiocarbamates.	Gopinath and Ford (64)	
Phase II				
Induction	Sodium sulfate and certain arylamines	Increased sulfate supply might result in a greater amount available for conjugation of xenobiotics and their metabolites.	—	
Inhibition	Pentachlorophenol and arylamines	Pentachlorophenol, an inhibitor of cytosolic sulfotransferases, renders such compounds as <i>N</i> -hydroxy-2-acetylaminofluorene (that are activated by the formation of sulfate esters) less toxic.	Meerman et al. (65,66)	
Excretion				
Pulmonary	Ethanol and mercury	Ethanol depresses the conversion of elemental mercury into ionic form. Their coexistence results in a diminution of the pulmonary retention as well as blood levels of mercury and enhances its pulmonary exhalation.	Nielsen-Kudsk (67) Magos et al. (68)	
Biliary	Arsenic and selenium Mercury and selenium	Arsenic increases the clearance of selenium from the liver into the bile. Selenium, on the other hand, inhibits the biliary excretion of mercury.	Lavender and Baumann (69)	
Urinary	Sodium bicarbonate (NaHCO <sub>3</sub> ) and fluoride	Alkalosis induced by NaHCO <sub>3</sub> causes a more rapid renal clearance of fluoride.	Reynolds et al. (70); Whitford and Pashley	



**Figure 9.** A preliminary schematic for considering data on chemical interactions for regulation. Abbreviations: Y, yes; N, no; UCF I/II, uncertainty factors.

eventually cause supra-additive neurotoxic effects even at exposure concentrations that do not exceed their respective TLVs. However, these toxicokinetic interactions involving mainly the biotransformation system appear to have a threshold level for each component.

Toxic interactions demonstrated in nonoccupational settings, i.e., the general environment, mainly involve nutritionally-important metals. In this regard, antagonistic effects on the absorption of toxic metals have often been reported to occur with higher intake of iron and zinc. Importantly, these interactions occur even when the dietary intake of one of these metals is insufficient, and therefore can be regarded as being more relevant for humans than most of the other chemical interactions.

Another interaction that belongs to this category is that of selenium and several carcinogens. A number of epidemiologic studies have reported an inverse relationship between the selenium status and the incidence rates of various types of cancer and cardiovascular diseases in humans (78–80). Recently, Yu et al. (81) have reported that the primary liver cancer incidence in people (in the Chinese province of Jiangsu) consuming selenium supplemented table salt was lower than that in the population not receiving selenium supplementation.

### Interactions Demonstrated in Humans at High Exposure Concentrations Only

Certain chemical interactions occur only at high doses or after unusual exposure scenarios (e.g., accidental exposures). An example of this category is the ketone-haloalkane interaction. The suspected supra-additive interaction between isopropanol and carbon tetrachloride occurred in two instances, both when the environmental concentrations exceeded the allowable exposure limits (52,53). It is understandable that when enzyme induc-

tion is the mechanism involved, there probably is a threshold for interaction. Particularly, this type of interaction can be expected to be insignificant at low exposure concentrations, where the rate-limiting factor of metabolism is hepatic blood flow and not the capacity of the enzyme system. On the other hand, the enzyme inducers, on coexposure, may become inhibitors of metabolism of other chemicals, the importance of such an interaction being determined by the affinity and relative roles of various isoforms of P450 involved in the metabolism of both chemicals.

### Interactions Demonstrated in Animal Studies at Low Doses but Potential for Occurrence in Humans Not Known

Several chemicals acting as inhibitors of hepatic and extrahepatic metabolism appear to have greater potential for interaction than inducers at low exposure concentrations. Apparently, the quantitative importance of the influence of an inhibitor on another chemical depends on the magnitude of the inhibition constant,  $K_i$ . Based on experimental and simulation studies, *trans*-1,2-dichloroethylene (DCE) has been suggested to be a potent suicide inhibitor of P450 (82). Therefore, metabolic inhibition of a variety of chemicals by DCE can be expected during combined exposures, even though direct evidence in humans is not yet available. Based on mechanistic considerations, uncompetitive and non-competitive inhibitions are also quantitatively more important than competitive inhibition. Benzene-toluene interaction in animals has been described to be the result of noncompetitive interaction (83), indicating that this interaction can occur at low exposure concentrations. Evidence for mutual inhibitory interaction between these chemicals has actually been obtained in an occupational exposure monitoring study (42). Another example belonging to this category is the toxicodynamic interaction between chlordecone and carbon tetrachloride demonstrated at low doses in animal studies (84).

### Interactions Demonstrated in Animal Studies at High Doses but Potential for Occurrence at Low Doses Not Known

Most of the chemical interaction studies conducted to date belong to this category. Typically, these studies have involved the administration of high doses of one or both chemicals by routes and scenarios often different from anticipated human exposures.

The potential of these interactions to occur during low-dose, chronic exposures is not known. Consideration of the number of combinations that need to be tested at low doses in chronic experiments, by at least one exposure route, emphasizes the need to utilize alternative methodologies, particularly predictive modeling strategy. This quantitative approach involves the integration of mechanistic factors of chemical disposition, interaction, and effect into a biologically based framework, that provides the basis for predicting the extent of interactions in untested exposure situations (72).

There is also the possibility that an interaction, not occurring in animals at high or low doses, may in fact occur in humans exposed to low concentrations. Based on mechanistic considerations, such observations would suggest differences in the mechanistic determinants of interaction between the two species (e.g., trans-species variation in the relative roles of different P450 isozymes).

The preceding analysis of the relevance and relative importance of experimental data on chemical interactions for humans covered only certain of the important aspects. For example, the influences of exposure routes and sequence of chemical exposures in the interaction studies are not explicitly taken into account. These factors are being considered in a weight-of-evidence approach formulated by the Environmental Criteria Assessment Office of the U.S. Environmental Protection Agency (85).

In summary, there is some evidence for the occurrence of chemical interactions in humans, particularly in the occupational environment and during accidental exposures. We need to investigate and utilize quantitative modeling approaches in the study of interactions to uncover the animal interactions that are relevant to human exposure situations. Such a quantitative, mechanistic approach to the study of chemical interactions is fundamentally important to achieve the ultimate goal of assessing the health risks associated with human exposure to complex chemical mixtures.

## REFERENCES

1. NAS. Principles of Toxicological Interactions Associated with Multiple Chemical Exposures. Washington:National Academy of Sciences, (1983).
2. Krishnan K, Brodeur J. Toxicological consequences of combined exposures to environmental pollutants. *Arch Complex Environ Studies* 3(3):1-106 (1991).
3. Waldbott GL. Pulmonary irritants. In: *Health Effects of Environmental Pollutants*. St. Louis:CV Mosby and Co, 1973;84.
4. Bates DV, Hazucha M. The effects of low levels of sulfur dioxide and ozone in the same atmosphere on the pulmonary function. In: *Proceedings of an International Symposium on the Recent Advances in the Assessment of the Health Effects of Environmental Pollution*, Vol 4. Geneva:World Health Organization, 1974;1977-1988.
5. Pershageni G, Wall S, Taube A, Limman L. On the interaction between occupational arsenic exposure and smoking and its relationship to lung cancer. *Scand J Work Environ Health* 7:302-309 (1981).
6. McJilton CE, Salawich RJ, Wofsy SC, Logan JA. Role of relative humidity in the synergistic effects of a sulfur dioxide aerosol mixture in the lung. *Science* 182:503-504 (1986).
7. Stacey RM, Seal EG, Housie AF, Green J, Roger LJ, Reggio L. Effects of gaseous and aerosol pollutants on pulmonary function measurement of normal humans. *Arch Environ Health* 38:104-115 (1983).
8. Bell K, Linn W, Hazucha M, Hackney J, Bates D. Respiratory effects of exposure to ozone plus sulfur dioxide in southern California vs eastern Canadians. *Am Ind Hyg Assoc J* 38:696-706 (1977).
9. Bedi JF, Folinsbee LJ, Horvath SM, Ebenstein RS. Human exposure to sulfur dioxide and ozone: absence of a synergistic effect. *Arch Environ Health* 34:233-239 (1979).
10. Horvath SM, Folinsbee LJ, Bedi JF. Combined effect of ozone and sulfuric acid on pulmonary function in man. *Am Ind Hyg Assoc J* 48:94-98 (1987).
11. Frank R. Sulfur dioxide-particulate interactions: recent observations. *Am J Ind Med* 1:427-434 (1980).
12. Baker EL, Zack M, Miles JW, Alderman L, Warren M, Dobbin RD, Miller S, Teekerr WR. Epidemic malathion poisoning in Pakistan malaria workers. *Lancet* 1: 31-38 (1978).
13. Aldridge WN, Norseth T. The toxicological properties of impurities in malathion. *Arch Toxicol* 42:95-106 (1979).
14. Miles JW, Mount DL, Staiger MA, Teeters WR. The S-methyl isomer content of stored malathion and fenitrothion water dispersible powders and its relationship to toxicity. *J Food Agri Chem* 27:421-425 (1979).
15. Parizek J, Ostadalo I, Kalouskova J, Babicky A, Benes J. The detoxifying effects of selenium interrelations between compounds of selenium and certain metals. In: *Newer Trace Elements in Nutrition* (Metz W, Cornatzer WE, eds). New York:Marcel Dekker, 1971;85-122.
16. Parizek J, Kalouskova J, Babicky A, Benes J, Pavlik L. Interaction of selenium with mercury, selenium, cadmium, and other toxic metals. In: *Trace Elements Metabolism in Animals*, Vol 2 (Hoekstra WG, Sutties TF, Ganther HE, Metz W, eds). Baltimore:University Park Press, 1974;119-131.
17. Ganther HE. Modifications of methyl mercury toxicity and metabolism by selenium and vitamin E: possible mechanisms. *Environ Health Perspect* 25:71-76 (1978).
18. Stopford W, Donovan DH, Abou-Donia MB, Menzel DB. Glutathione peroxidase deficiency and mercury allergy: amelioration with selenium supplementation. In: *Proceedings of the Symposium on Selenium-Tellurium in the Environment*. Pittsburgh, PA:Industrial Health Foundation, 1976;105-112.
19. Suzuki T, Satoh H, Yamamoto R, Kashiwarzaki H. Selenium and mercury in food stuff from a locality with elevated intake of mercury. *Bull Environ Contam Toxicol* 24:805-811 (1980).
20. Kostaf L, Brynet AR, Zelenko V. Correlation between selenium and mercury in man following exposure to inorganic mercury. *Nature (London)* 254:238 (1975).
21. Nylander M, Weiner J. Mercury, selenium concentrations and their interrelationships in the organs from dental staff and the general population. *Br J Ind Med* 48:729-734 (1991).
22. Barber TE, Menke RD. The relationship of ingested iron to the absorption of mercuric oxide. *Am J Emerg Med* 2:500-505 (1984).
23. Six KM, Goyer RA. The influence of iron deficiency on tissue content and toxicity of ingested lead in the rat. *J Lab Clin Med* 79:128-136 (1972).
24. Ragan HA. Effects of iron deficiency on absorption and distribution of lead and cadmium in rats. *J Lab Clin Med* 90: 700-706 (1977).
25. Barton JC, Conrad ME, Nuby S, Harrison L. Effects of iron on the absorption and retention of lead. *J Lab Clin Med* 92:937-947 (1978).
26. Flanagan PR, Hamilton DL, Haist J, Valberg L. Interrelationships between iron and lead absorption in iron-deficient mice. *Gastroenterology* 77:1074-1081 (1979).
27. Dhir H, Sharma A, Talikar G. Alteration of cytotoxic effects of lead through interaction with other heavy metals. *Nucleus* 28:69-89 (1985).
28. Mahaffey KR. Nutritional factors and susceptibility to lead toxicity. *Environ Health Perspect* 8:107-112 (1974).
29. Sullivan JF, Reummler PS. Effect of excess iron on cadmium or lead absorption in rats. *J Toxicol Environ Health* 22:131-139 (1987).
30. Ito Y, Niiya Y, Otani M, Sarai S, Shima S. Effect of food intake on blood lead concentration in workers occupationally exposed to lead. *Toxicol Lett* 37:105-114 (1987).
31. Chisolm JJ. Dose-effect relationships for lead in young children: evidence in children for interactions among zinc and iron. In: *Environmental Lead: Proceedings of the Second International Symposium on Lead Research* (Lynam DR, Piantandia LG, Cole LG, eds). New York:Academic Press, 1981;1-7.
32. Watson WS, Hume R, Moore MR. Oral absorption of lead and iron. *Lancet* 2: 236-237 (1980).
33. Yip R, Norris TN, Anderson AS. Iron status of children with elevated blood lead concentrations. *J Pediatr* 98:922-925 (1981).
34. Mahaffey KR, Annett JL. Assessment of erythrocyte protoporphyrin with blood lead level and iron status in the Second National Health and Nutrition examination survey 1976-1980. *Environ Res* 41:327-336 (1986).
35. Dukiewicz B, Dukiewicz T, Milkowska G. The effect of mixed exposure to lead and zinc on aminolevulinic acid level in urine. *Int Arch Occup Environ Health* 42:341-348 (1979).
36. Papaioannou R, Sohler A, Pfiffer CC. Reduction of blood lead levels in battery workers by zinc and vitamin C. *J Orthomol Psychiatry* 7: 1-13 (1978).
37. Flanagan PR, Chamberlain MJ, Valberg LS. The relationship between iron and lead absorption in humans. *Am J Clin Nutr* 36:823-829 (1982).
38. Lauwerys R, Roelofs H, Buchet JP, Bernard AA, Verhoeven L, Konings L. The influence of orally administered vitamin C or zinc on the absorption of and biological response to lead. *J Occup Med* 25:668-674 (1983).
39. Solomons NW. Competitive interaction of iron and zinc in the diet: consequences for human nutrition. *J Nutr* 116:927-935 (1986).
40. Bunker VW, Lawson MS, Delves HT, Clayton BE. The intake and excretion of lead and cadmium by the elderly. *Am J Clin Nutr* 39:803 (1984).
41. Pfaffli P, Backman AL. Trichloroethylene concentrations in blood and expired air as indicators of occupational exposure: a preliminary report. *Scand J Work Environ Health* 9:140-144 (1972).

42. Inoue O, Seiji K, Watanabe T, Kasahara M, Nakatsuka H, Yin S, Jin C, Ikeda M. Mutual metabolic suppression between benzene and toluene in man. *Int Arch Occup Environ Health* 60:15–20 (1988).
43. Angerer J, Lehnert G. Occupational chronic exposure to organic solvents. 8. Phenolic compounds—metabolites of alkylbenzenes in man: simultaneous exposure to ethylbenzene and xylenes. *Int Arch Occup Environ Health* 43:145–150 (1979).
44. Tardif R, Lapare S, Plaa GL, Brodeur J. Effect of simultaneous exposure to toluene and xylene on their respective biological exposure indices in humans. *Int Arch Occup Environ Health* 63:279–284 (1991).
45. Inoue O, Seiji K, Kawai T, Jin YT, Cai SX, Yin SN, Li GL, Nakatsuka H, Watanabe H, Ikeda M. Relationship between vapor pressure and urinary metabolite excretion among workers exposed to trichloroethylene. *Am J Ind Med* 15:103–110 (1988).
46. Liira J, Riihimäki V, Engstrom K. Coexposure of man to m-xylene and methyl ethyl ketone. *Scand J Work Environ Health* 14:322–327 (1988).
47. Riihimäki V. Percutaneous absorption of m-xylene from a mixture of m-xylene and isobutanol in man. *Scand J Work Environ Health* 5:143–150 (1979).
48. Olson BA, Gamberale F, Iregren A. Co-exposure to toluene and p-xylene in man: central nervous systems. *Br J Ind Med*, 42:117–122 (1985).
49. Yin S, Li G, Hu Y, Zhang X, Jin C, Inoue O, Seiji K, Kasahara M, Nakatsuka H, Ikeda M. Symptoms and signs of workers exposed to benzene, toluene, or their combination. *Ind Health* 25:113–130 (1987).
50. Pass MA, Muir AD, Majak W, Yost G. Effect of alcohol and acetaldehyde dehydrogenase inhibitions on the toxicity of 3-nitropropanol in rats. *Toxicol Appl Pharmacol* 78:310–315 (1985).
51. Brady JF, Li MJ, Ishizaki H, Lee M, Ning S, Xiao F, Yang CS. Induction of cytochrome P-450IIE1 by secondary ketones and the role of P-450IIE1 in chloroform metabolism. *Toxicol Appl Pharmacol* 100:342–349.
52. Folland DS, Schaffner W, Grinn H, Crofford QB, McMurray DK. Carbon tetrachloride potentiation by isopropanol: investigation of an industrial outbreak. *J Am Med Assoc* 236:1853–1856 (1976).
53. Deng JF, Wang JD, Sih TS, Lan FL. Outbreak of carbon tetrachloride poisoning in a color printing factory related to the use of isopropyl alcohol and air conditioning system in Taiwan. *Am J Ind Med* 12:11–19 (1987).
54. Allen N, Mendell JR, Billmaier D, Fontaine RE, Oeill J. Toxic polyneuropathy due to methyl n-butyl ketone. *Arch Neurol* 32:209–218 (1975).
55. Billmaier D, Yee HT, Allen N, Craft B, Williams N, Epstein S, Fontaine R. Peripheral neuropathy in a coated fabrics plant. *J Occup Med* 16: 665–671 (1974).
56. Altenkirch H. Schnuffelsucht Chronischer Lösungsmittel Missbrauch bei Kindern und Jugendlichen mit neurotoxischen Folgen. *Dtsch Med Wochenschr* 104:935–938 (1979).
57. Jacob SW, Bischel M, Herschler RJ. Dimethyl sulfoxide: effects on the permeability of biologic membranes. *Curr Ther Res* 6:193 (1964).
58. Hayes WJ, Pearce GW. Pesticide formulation: relation to safety in use. *J Agr Food Chem* 1:466 (1953).
59. Dreisbach RH. Cyanides, sulfides and cobalt. In: *Handbook of Poisoning*. 9 ed. Los Altos, CA: Lange Medical Publishing Co, 1977;241.
60. Conrad ME, Barton JC. Factors affecting the absorption and excretion of lead. *Gastroenterology* 74:731–740.
61. Osakrsson A, Lind B. Increased lead levels in brain after long term treatment with lead and dithiocarbamate or thium derivatives in rats. *Acta Pharmacol Toxicol* 56: 309–315 (1985).
62. Triolo AJ, Coon JM. Binding of paraoxon by plasma of aldrin-treated mice. *Toxicol Appl Pharmacol* 14:622 (1970).
63. Cohen SD, Murphy SD. A simplified bioassay for organophosphate detoxication and interactions. *Toxicol Appl Pharmacol* 27:537–550.
64. Gopinath C, Ford EJH. The role of microsomal hydroxylases in the modification of chloroform hepatotoxicity in the rat. *Br J Exp Pathol* 56:412–422 (1975).
65. Meerman JHN, van Doorn ABD, Mulder GJ. Inhibition of sulfate conjugation of N-hydroxy 2-acetylfluorene in isolated perfused rat liver and in the rat *in vivo* by pentachlorophenol. *Cancer Res* 40: 3772–3779 (1980).
66. Meerman JHN, Sterenborg D, Mulder GJ. Use of pentachlorophenol as longterm inhibitor of sulfation of phenols and hydroxamic acids in the rat liver *in vivo*. *Biochem Pharmacol* 32:1587–1593 (1983).
67. Nielsen-Kudsk F. The influence of alcohol in the absorption of mercury vapor from the lungs in man. *Acta Pharmacol Toxicol* 23:263–274.
68. Magos L, Clarkson TW, Greenwood MR. The depression of pulmonary retention of mercury vapor by ethanol: identification of the site of action. *Toxicol Appl Pharmacol* 26:180–183 (1973).
69. Lavender OA, Baumann CA. Selenium metabolism: effect of arsenic on the excretion of selenium in the bile. *Toxicol Appl Pharmacol* 9: 106–115 (1966).
70. Reynolds KE, Whitford GM, Pashley DH. Acute fluoride toxicity: the influence of acid-base status. *Toxicol Appl Pharmacol* 45:415–427 (1978).
71. Whitford GM, Pashley DH. Fluoride reabsorption by nonionic diffusion in distal nephron of the dog. *Proc Soc Exp Biol Med* 196:178 (1991).
72. Krishnan K, Clewell HJ III, Andersen ME. Physiologically based pharmacokinetic modeling of simple mixtures. *Environ Health Perspect* 102(Suppl 9):151–155 (1994).
73. Frawley JP, Fuyat HN, Hagan EC, Blake JR, Fitzugh OG. Marked potentiation of mammalian toxicity from simultaneous administration of two cholinesterase compounds. *J Pharmacol Exp Ther* 121:96–106 (1957).
74. DuBois KP. Combined effects of pesticides. *Can Med Assoc J* 100:173–179 (1968).
75. Moeller HC, Clary JJ. Potentiation of red blood cell cholinesterase activity as indicators of the threshold of incipient toxicity of EPN and malathion in human beings. *Toxicol Appl Pharmacol* 4:123–130 (1962).
76. Last JA. Toxicological interactions between inorganic gases and particulates. *Fundam Appl Toxicol* 12:9–11 (1989).
77. Last JA. Synergistic effects of air pollution: ozone plus a respiratory aerosol. Research Report No. 38. Cambridge, MA: Health Effects Institute, 43, 1990.
78. Schrauzer GN, White DA, Schneider CJ. Cancer mortality studies 3: statistical association with dietary selenium intakes. *Bioinorg Chem* 7:23–34 (1977).
79. Shamberger RJ, Tyoko SA, Willis CA. Antioxidant and cancer: 4. Selenium and age adjusted mortality. *Arch Environ Health* 31:231–235 (1976).
80. Kok FJ, Hofman A, Witteman JCM, De Bruin AM, Kryossen DCM, De Bruin M, Valkenburg HA. Decreased selenium levels in acute myocardial infarction. *J Am Med Assoc* 261:1161–1164 (1989).
81. Yu SY, Zhu YJ, Li WG, Hou C. Chemoprevention trial of primary liver cancer with selenium supplementation in Qidong county of China. In: *Metal Ions in Biology and Medicine* (Collery PH, Poirier LA, Manfait JC, Etienne JC, eds). Paris: John Libbey Eurotext, 1990;497–499.
82. Andersen ME, Gargas ML, Clewell HJ III. Suicide inactivation of microsomal oxidation of cis- and trans- dichloroethylene in male Fischer-344 rats *in vivo*. *Toxicologist* 6:12 (1986).
83. Purcell KJ, Cason GH, Gargas ML, Andersen ME, Travis CC. *In vivo* metabolic interactions of benzene and toluene. *Toxicol Lett*, 52:141–152 (1990).
84. Mehendale HM. Mechanism of lethal interaction of chloroform and carbon tetrachloride at nontoxic doses. *Toxicol Lett* 49: 215–241 (1989).
85. Mumtaz MM. The US EPA guidelines and current issues of the health risk assessment of chemical mixtures. *Toxicologist* 12:4 (1992).